

OSURG Prophylactic Antibiotics (Regimen for Selected Surgeries)

<p style="text-align: center;"><u>Pacemakers or AICDs or Vascular</u></p> <p style="text-align: center;">Cefazolin, Cefuroxime, or Vancomycin**</p> <p style="text-align: center;">If beta-lactam allergy: Vancomycin* or Clindamycin*</p>	<p style="text-align: center;"><u>Orthopedic/Podiatry</u></p> <p style="text-align: center;">Cefazolin, Cefuroxime, or Vancomycin**</p> <p style="text-align: center;">If beta-lactam allergy: Vancomycin* or Clindamycin*</p>
<p style="text-align: center;"><u>Genitourinary – Prostate Biopsy</u></p> <p style="text-align: center;">Quinolone (Ciprofloxacin) Or 1st Generation cephalosporin (Cefazolin) Or Aminoglycoside (Gentamicin) AND Metronidazole (Flagyl) Or Aminoglycoside (Gentamicin) AND Clindamycin</p> <p style="text-align: center;"><u>Genitourinary – Penile Prosthesis insertion, removal, revision</u></p> <p style="text-align: center;">Ampicillin/Sulbactam Or Aminoglycoside (Gentamicin) AND 1st Generation cephalosporin (Cefazolin)</p> <p style="text-align: center;">If beta-lactam allergy: Aminoglycoside (Gentamicin) AND Vancomycin</p>	<p style="text-align: center;"><u>Hysterectomy</u></p> <p style="text-align: center;">Cefazolin or Cefoxitin</p> <p style="text-align: center;">If beta-lactam allergy: Clindamycin AND Gentamicin Or Metronidazole AND Ciprofloxacin</p> <p style="text-align: center;"><u>Pubovaginal Sling</u></p> <p style="text-align: center;">Cefazolin, Cefotetan, or Cefoxitin</p> <p style="text-align: center;">If beta-lactam allergy: Gentamicin AND Metronidazole</p>
<p style="text-align: center;"><u>PEG Placement</u></p> <p style="text-align: center;">Cefazolin</p> <p style="text-align: center;">If beta-lactam allergy: Gentamicin AND Vancomycin</p>	<p style="text-align: center;"><u>Head and Neck Procedures</u></p> <p style="text-align: center;">Cefazolin or Vancomycin**</p> <p style="text-align: center;"><u>Neurological Procedures</u></p> <p style="text-align: center;">Cefazolin</p> <p style="text-align: center;">If beta-lactam allergy: Vancomycin or Clindamycin</p>

*For cardiac, orthopedic, and vascular surgery, if the patient is allergic to beta-lactam antibiotics, Vancomycin or Clindamycin are acceptable substitutes

**Vancomycin is acceptable with a Physician/APN/PA/Pharmacist documented justification for its use

Special Note: Utilize the specific SCIP surgery prophylactic antibiotic order forms located on the Meridian Intranet under Clinical Forms. These forms provide the most current recommended antibiotics and antibiotics available on formulary.

